

# Impact of small ruminant diseases on different household members in Ethiopia

Barbara Wieland, Biruk Alemu, Hiwot Desta, Wole Kinati, Shiferaw Tafesse, Anouka Van Eerdewijk, Annet A. Mulema

First joint conference of the Association of Institutions for Tropical Veterinary Medicine and the Society of Tropical Veterinary Medicine, Berlin, Germany, 4–8 September 2016



# Background

- Small ruminants serve multiple livelihood roles for Ethiopian small holders
- Women play important role in small ruminant production
- Little is known on how disease impact and risk mitigation strategies differ among household members

→ understanding of these issues is pivotal to design sustainable livestock health interventions

# Objectives

This study aimed at

- identifying disease constraints
- assess impacts of disease constraints as perceived by men and women in mixed crop-livestock and agro-pastoralist systems

# Materials and methods

## 1. Focus groups discussions (FGDs)

- In 23 villages in 4 regions, Oromia, Tigray, Amhara and SNNPR, in Ethiopia
- In each village separate FGDs were held with women, men, young women and young men
- Participatory tools: simple ranking, proportional piling and seasonal calendar



# Materials and methods

## 2. Household survey

- 440 household in the same areas
- 50% of interviews conducted with women, 50% with men
- Questions based on FGD findings



# Quantify/rank impacts

## In FGDs:

For identified important diseases, farmers were asked

- how the disease impacts the household
- which household members most affected
- proportional piling for M, W, YM, YF, CH

## In HH survey:

- 3 main impacts for the 3 most important diseases
- recorded according to impact categories from FGDs



# Results


## Disease Priority: agro-ecology

Disease Category	Highland Woredas		Lowland Woredas	
	Mean rank score	Test statistics	Mean rank score	Test statistics
Respiratory dis	<u>6.10</u>	N=68	4.40	N=24
Neurological dis	4.40	Kendall's W= <u>0.38</u>	3.23	Kendall's W= <u>0.09</u>
Skin dis	3.93	Chi-Square= 153.952	4.15	Chi-Square=12.382
GIT	<u>5.01</u>	Df=6	4.44	Df=6
External par	2.71	Asymp. Sig.= 0.000	3.69	Asymp. Sig.= <u>0.054</u>
Systemic	2.72		<u>4.81</u>	
Other	3.13		<u>3.29</u>	

- In highland areas (mixed crop-livestock system): priorities were respiratory diseases and GIT parasites with strong agreement among respondent groups
- In lowland areas (pastoral and agro-pastoral system): priorities were systemic diseases and neurological diseases, differences between regions

# Results

## Disease priority: gender



- Women and young female scored respiratory diseases higher than young male and men
- Young men and women scored neurological diseases higher than men and young female



# Results

## Impact categories identified in FGDs

### Economical


- Financial: loss of income
- Mortality
- Productivity: poor growth rates & weight gain, lack of milk for offspring
- Value: meat quality, hide quality
- Costs: for treatment
- Waste of time

### Social

- Social/psychological: status, taxes unpaid, no mixing with other animals
- Drop out of school: school fees unpaid
- Migration for other jobs in cities

### Human

- Food security/malnutrition
- Human health



■ Financial /Loss of income

■ Mortality Loss of animal/herd

■ Low/loss of productivity

■ Additional unexpected cost (Sell other animal or cereal crop)

■ Wastage of time treating the animals

■ Migration for other jobs

■ Drop out of school

■ psychological impact

■ Social disturbance/ isolation


■ Malnutrition

■ Human health

# Who is affected from the HH? Why?


Women are most severely affected

- women bear the main responsibility of looking after diseased animals
- In order to fulfill their household duties (children, food), they rely on income from small ruminants
- Lack of other income sources if animals are lost due to disease
- For men it is easier find other work if for some reason animals are lost




Kendall's W = 0.49, p=0.000  
Error bars: 95% CI


# Impact by disease categories



# Impact perception: by gender



# Impact: Agro-ecology



# Conclusions

- Impact of SR diseases highly important in women
- Understanding of impacts almost similar in men and women
- Need to take gender into account when designing small ruminant health interventions



# Acknowledgements

This work is financed by

- SmarT project (IFAD), Africa RISING project (USAID)

It is implemented in a partnership with

- ICARDA, Regional Research Centers (DBARC, SDARC, YPDARC, BongaARC, AbergelleARC, BakoARC, ArekaARC), Mada Walabu University and Agricultural Offices in the project sites

It contributes to the CGIAR Research Program on

- Livestock and Fish

*better lives through livestock*

ilri.org

ilri.org  
*better lives through livestock*  
ILRI is a member of the CGIAR Consortium

Box 30709, Nairobi 00100 Kenya  
Phone +254 20 422 3000  
Fax +254 20 4223001  
Email [ilri-kenya@cgiar.org](mailto:ilri-kenya@cgiar.org)

ILRI has offices in:  
Central America • East Africa  
• South Asia • Southeast and East Asia  
• Southern Africa • West Africa



The presentation has a Creative Commons license. You are free to re-use or distribute this work, provided credit is given to ILRI.